

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 March 2001 (08.03.2001)

PCT

(10) International Publication Number
WO 01/15688 A1

(51) International Patent Classification⁷: A61K 31/19, 9/48

(21) International Application Number: PCT/US00/23908

(22) International Filing Date:
1 September 2000 (01.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/389,003 2 September 1999 (02.09.1999) US

(71) Applicant: BANNER PHARMACAPS, INC. [US/US];
4125 Premier Drive, High Point, NC 27261-2210 (US).

(72) Inventor: GULLAPALLI, Rampurna, Prasad; 1921
New Garden Road, Apt. N-8, Greensboro, NC 27410
(US).

(74) Agent: HICKS, Jack, B.; Rhodes & Mason, P.L.L.C., P.O.
Box 2974, Greensboro, NC 27402 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: IBUPROFEN-CONTAINING SOFTGELS

(57) Abstract: Liquid softgel fill formulations containing ibuprofen in free acid form, and softgel capsules comprised of a gelatin sheath enclosing such fill formulations, are prepared by dissolving more than 30 % of ibuprofen in free acid form in polyethylene glycol and at least 10 % by weight of a polyvinylpyrrolidone having an average molecular weight of from about 2,000 to about 54,000. The formulations may also include a surfactant to increase the bioavailability of the ibuprofen.

WO 01/15688 A1

IBUPROFEN-CONTAINING SOFTGELS

Background of the Invention

5 (1) Field of the Invention

The present invention relates generally to softgels or soft gelatin capsules and fill compositions therefor, and in particular to a soft gelatin capsule that contains a highly concentrated solution of ibuprofen in its free form.

10 (2) Description of the Prior Art

U.S. Patent No. 4,690,823 to Lohner et al. describes a process for manufacturing ibuprofen-containing soft gelatin capsules in which up to 30 parts by weight of ibuprofen in free form is dissolved in from 70 to 85 parts by weight of polyoxyethylene-polyoxypropylene polymer or in a mixture of from 30 to 76 parts by weight of a
15 polyalkylene glycol and from 7 to 40 parts by weight of a surfactant having a very rapid and high bio-availability of the active ingredient. Preferred suitable surfactants include, for example, polyoxyethyleneglycerol trihydroxystearate, polyoxyethylene (C₁₂₋₁₈)-fatty alcohol ethers, polyoxyethylene stearate, polyoxyethylenesorbitan mono(C₁₂₋₁₈)-fatty acid esters, and polyoxyethylene-polyoxypropylene polymer.

20 In the Lohner et al. process, up to 30 parts by weight of free form ibuprofen is dissolved by heating the free form ibuprofen with the selected solvent at a temperature of 45° to 60° C. If a concentration of ibuprofen greater than the maximum 30 parts by weight possible in solution is desired, up to 40 parts additional parts by weight of ibuprofen may also be suspended in the fill.

Softgel fills containing high concentrations of ibuprofen as a mixture of ibuprofen in free form and its salts are described in U.S. Patent Nos. 5,071,643 and 5,360,615 to Yu et al. These fill formulations may be liquid, semi-solid or solid, and are formed by mixing 40-80% by weight ibuprofen, 0.1-1.5 moles of hydroxide ion per mole of
5 ibuprofen, 1-20% by weight water, and 4-12% by weight glycerine or propylene glycol in polyethylene glycol. Solubility of the ibuprofen salts is further enhanced 2-10% by the further addition of 3-10% by weight of glycerin, or propylene glycol or 1-20% by weight of polyvinylpyrrolidone.

The preferred average molecular weight of the polyvinylpyrrolidone is 10,000-
10 100,000. Higher percentages of above 5% polyvinylpyrrolidone, and higher molecular weight polyvinylpyrrolidone are used to prepare semi-solid and solid formulations for suppositories, two piece capsules, and tablets.

Upon mixing under the conditions described by Yu et al., the polyethylene glycol acts to dissolve the free form of the ibuprofen; the hydroxyl ions source, e.g., sodium
15 hydroxide or potassium hydroxide, partially forms an ibuprofen salt, and the water forms a solvation sphere around the acid salt permitting it to go into solution in the polyethylene glycol.

While softgel fill compositions containing ibuprofen salts can be produced by the above procedures, there is still a need for liquid softgel fill formulations with high
20 concentrations, i.e., greater than 30% by weight, of ibuprofen in solution in its free acid form. In particular, there is a need for highly concentrated solutions containing ibuprofen

in its free form that do not adversely affect the softgel capsules over an extended period of time.

Summary of the Invention

The present invention relates to a liquid softgel fill formulation comprising greater
5 than 30 % by weight ibuprofen in free acid form in solution; from about 30 to about 60%
by weight polyethylene glycol; and from about 10 to about 30 % by weight of
polyvinylpyrrolidone.

More specifically, liquid softgel fill formulations containing ibuprofen in free acid
form, and softgel capsules comprised of a gelatin sheath enclosing such fill formulations,
10 are prepared by dissolving more than 30% of ibuprofen in free acid form in polyethylene
glycol and at least 10% by weight of a polyvinylpyrrolidone having an average molecular
weight of from about 2,000 to about 54,000. The formulations may also include a
surfactant to increase the bioavailability of the ibuprofen.

Detailed Description of The Invention

15 The present invention relates to softgel fill formulations containing high
concentrations, i.e., greater than 30% by weight of ibuprofen in its free acid form, and in
particular to highly concentrated, preferably non-aqueous, ibuprofen fill formulations.
Such formulations, when encapsulated into soft gelatin one-piece capsule sheaths by
known techniques, provide long term stability, while still being of an acceptable size to
20 the consumer. When used in the following description, "ibuprofen" will be understood to
mean ibuprofen in its free acid form.

It has been found that these formulations can be prepared by dissolving ibuprofen in a solvent comprised of polyethylene glycol and polyvinylpyrrolidone, with each of the solvent components being present in a defined amount in the composition, and with each component having defined characteristics. Preferably, the formulation also includes a
5 surfactant to enhance bioavailability of the ibuprofen.

Specifically, the formulations of the present invention are comprised of greater than 30% and up to 40% or more by weight ibuprofen, from 30 to 60 % by weight polyethylene glycol, and from 10 to 30 % by weight of polyvinylpyrrolidone. Preferably, the formulations are comprised of from 35 to 40% by weight ibuprofen, from 30 to 50 %
10 by weight polyethylene glycol, and from 15 to 30% by weight of polyvinylpyrrolidone.

It will be understood that the percentages of the above ingredients in a given formulation, which may also have other ingredients present, will be adjusted to total 100%. For example, up to 6% of a nasal decongestant, such as pseudoephedrine hydrochloride, or up to 10% of an antihistaminic, such as diphenhydramine
15 hydrochloride, may be included in the formulation.

The preferred embodiments of the invention also contain at least about 10%, preferably from about 1% to about 10% by weight of a surfactant. Preferably, the surfactants are characterized by the capability to enhance the bioavailability of the ibuprofen, while being miscible with polyethylene glycol to form a clear solution.
20 Suitable surfactants include esters of d-alpha tocopheryl, polyoxyethylene castor oil derivatives, and polyglycolized glycerides. Poxoxyethylene castor oil derivatives are the reaction products of ethylene oxide and castor oil or hydrogenated castor oil sold, for

example, as Cremophors by BASF. Suitable polyglycolized glycerides are sold under the trademark Gelucire by Etablissements Gattefosse. An especially preferred surfactant is d-alpha tocopheryl polyethylene glycol 1000 succinate sold by Eastman Chemical Company under the trademark Vitamin E TPGS.

5 The polyethylene glycol component of the formulation should have an average molecular weight of up to about 1,000 in order to provide a liquid fill formulation. Preferably, the polyethylene glycol has an average molecular weight of at least about 200, with an average molecular weight of from about 400 to about 1,000 being preferred. PEG 600 is especially preferred.

10 The average molecular weight of the polyvinylpyrrolidone used in the present formulations may be in the range of from 2,000 to 54,000, but preferably is in the range of from about 2,000 to about 10,000. Preferred polyvinylpyrrolidones are sold under the trademarks Kollidon 12 PF and Kollidon 17 PF by BASF. These lower molecular weight polyvinylpyrrolidones have approximate molecular weights of 2,500 and 10,000
15 respectively. Another suitable polyvinylpyrrolidone is sold under the trademark Plasdane C-15 by ISP, and has a molecular weight of about 8,000. Unlike Yu, the present invention employs larger amounts of lower molecular weight polyvinylpyrrolidones to ensure that a liquid fill is obtained.

 An additional advantage of these higher amounts of polyvinylpyrrolidone is the
20 apparent reduction in the esterification reaction between the ibuprofen and the polyethylene glycol, a known disadvantage of mixing these two ingredients that results in the reduction of available ibuprofen in its free acid form. Use of the higher

percentages of polyvinylpyrrolidone permits a corresponding reduction in the amount of polyethylene glycol required.

The fill formulations may be prepared by heating the polyethylene glycol and surfactant to $55 \pm 5^\circ\text{C}$, while mixing. The ibuprofen and polyvinylpyrrolidone are added
5 and dissolved during heating and mixing. The procedure is carried out under a nitrogen atmosphere, and the final formulation is deaerated.

The fill formulation is encapsulated into one-piece gelatin sheath or shell that includes a plasticizer to control the softness and flexibility of the sheath, water, and optionally, other additives, such as flavorants, colorants, opacifiers, etc. The softgel
10 capsules may be produced in a known manner with a rotary die process in which a molten mass of a gelatin sheath formulation is fed from a reservoir onto drums to form two spaced sheets or ribbons of gelatin in a semi-molten state. These ribbons are fed around rollers and brought together at a convergent angle into the nip of a pair of roller dies that include opposed die cavities. A fill formulation to be encapsulated is fed into the wedge-
15 shaped joinder of the ribbons.

The gelatin ribbons are continuously conveyed between the dies, with portions of the fill formulation being trapped between the sheets inside the die cavities. The sheets are then pressed together, and severed around each die so that opposed edges of the sheets flow together to form a continuous gelatin sheath around the entrapped
20 medicament. The part of the gelatin sheet that is severed from the segments forming the capsules is then collected for recycling, and the soft capsules are dried.

Various sheath formulations known in the prior art may be used to encapsulate the fill formulations of the present invention. For example, suitable sheath formulations may include from about 35 to about 50 % by weight gelatin; at least 20% by weight, and preferably up to about 40 % by weight, of a plasticizer; and from about 25 to about 50 %
5 by weight water. These formulations, when formed into capsules and dried, will result in capsule sheaths comprised of from about 45 to about 75 % by weight gelatin; from about 20% to about 40 % by weight plasticizer; and from about 5 to about 15 % by weight water.

The gelatin will normally have a bloom in the range of from about 150 to about
10 275, and may be Type A or B gelatins or a mixture thereof. Limed bone, acid bone, fish and/or pig skin gelatins may be used.

The sheath plasticizer preferably is sorbitol, sorbitol special (a mixture of sorbitol and sorbitan), maltitol, or a mixture thereof. While glycerin can be used as a plasticizer, it has been found that the ibuprofen may esterify with the glycerin, reducing the amount
15 of available free form ibuprofen. Therefore, the non-glycerin plasticizers are preferred.

The sheath formulations may also contain other ingredients, such as taste modifiers, coloring agents, and moisture retaining agents. Taste modifiers include non-reducing sugars, such as xylitol, maltitol, or Lycasin[®] manufactured by Roquette America, Inc. of Keokuk, IA and normally will comprise up to about 5% by weight of the
20 sheath composition. Suitable moisture retaining agents include celluloses, cellulose derivatives, starches, starch derivatives, vegetable gums, non-hygroscopic, mono-, di-

and oligosaccharides, and silicon dioxide. Various FD&C coloring agents may be used to impart the desired color to the capsule.

In order to be acceptable to the consumer, the softgel capsule should be of a size that is easily swallowed. Generally, the fill size of the capsule will be less than 600 mg, and preferably about 500 mg or less, for the capsule to be of an acceptably small dimension. The required effective dosage of ibuprofen will normally be at least 175 mg, and preferably about 200 mg. Therefore, the preferred capsules will contain at least about 35% by weight of ibuprofen.

The overall volume of the fill is also made possible by the fact that only polyethylene glycol and polyvinylpyrrolidone are required for dissolution, with a surfactant preferably being included to enhance bioavailability of the ibuprofen when ingested. Thus, the compositions may be free of water and other ingredients that increase the fill volume.

The following examples are for the purpose of illustrating the present fill formulation, and softgels comprised of these fill formulation, and are not to be taken as in any way limiting the scope of the invention. Examples 1-10 are representative fill formulations containing in excess of 30% by wt. ibuprofen in its free acid form. Examples 11-13 are representative sheath formulations used to encapsulate the fill formulations of Examples 1-10.

Example 1

Fill Ingredients	% by wt.
Ibuprofen	35
PEG 600	37
Kollidon 17 PF	22
Vitamin E TPGS	6

Example 2

Fill Ingredients	% by wt.
Ibuprofen	35
PEG 600	35
Kollidon 17 PF	20
Vitamin E TPGS	5
Water, purified	5

5

Example 3

Fill Ingredients	% by wt.
Ibuprofen	37
PEG 600	37
Kollidon 17	20
Vitamin E TPGS	6

Example 4

Fill Ingredients	% by wt.
Ibuprofen	35
PEG 600	40
Kollidon 30	15
Cremophor RH 40	5
Propylene glycol	5

10

Example 5

Fill Ingredients	% by wt.
Ibuprofen	35
PEG 600	33
Kollidon 17 PF	22
Gelucire 44/14	10

Example 6

Fill Ingredients	% by wt.
Ibuprofen	35
PEG 600	40
Kollidon 30	15
Cremophor RH 60	5

Propylene glycol	5
------------------	---

Example 7

Fill Ingredients	% by wt.
Ibuprofen	35
PEG 600	35
Kollidon 30	15
Cremophor RH 60	10
Propylene glycol	5

5

Example 8

Fill Ingredients	% by wt.
Ibuprofen	40
PEG 600	40
Kollidon 17 PF	20

10

Example 9

Fill Ingredients	% by wt.
Ibuprofen	40
PEG 600	40
Kollidon 17 PF	15
Vitamin E TPGS	5

Example 10

Fill Ingredients	% by wt.
Ibuprofen	40
PEG 600	30
Kollidon 17 PF	25

Vitamin E TPGS	5
----------------	---

Example 11

Sheath Ingredients	% by wt.
Gelatin, lime bone	42
Sorbitol, special	9
Maltitol, 75%	12
Water, purified	37

5

Example 12

Sheath Ingredients	% by wt.
Gelatin, lime bone	42
Sorbitol, special	15
Maltitol, 75%	6
Water, purified	37

Example 13

Sheath Ingredients	% by wt.
Gelatin, lime bone	42
Glycerin	8
Maltitol, 75%	12
Water, purified	38

10 Certain modifications and improvements will occur to those skilled in the art upon
 a reading of the foregoing description. It should be understood that all such
 modifications and improvements have been deleted herein for the sake of conciseness and
 readability but are properly within the scope of the follow claims.

What is claimed is:

1. A liquid softgel fill formulation comprising:

a) greater than 30 % by weight ibuprofen in free acid form in solution;

b) from about 30 to about 60% by weight polyethylene glycol; and

5 c) from about 10 to about 30 % by weight of polyvinylpyrrolidone.

2. The softgel fill formulation of claim 1, wherein said ibuprofen is present in an amount of at least about 35% by weight.

3. The softgel formulation of claim 1, wherein said polyethylene glycol has an average molecular weight of from about 200 to about 1,000.

10 4. The softgel formulation of claim 1, wherein said polyvinylpyrrolidone has an average molecular weight of from about 2,000 to about 54,000.

5. The softgel fill formulation of claim 1, further including up to about 6% of a nasal decongestant or up to about 10% of an antihistaminic.

15 6. The softgel fill formulation of claim 1, further including up to about 10% by weight of a surfactant to increase the bioavailability of said ibuprofen.

7. The softgel formulation of claim 6, wherein said surfactant is selected from the group consisting of esters of d-alpha-tocopheryl, polyoxyethylene castor oil derivatives, and polyglycolized glycerides.

8. A liquid softgel fill formulation comprising:

20 a) greater than 30% by weight ibuprofen in free acid form in solution;

b) from about 30 to about 60% by weight of polyethylene glycol having an average molecular weight of from about 200 to about 1,000;

c) from about 10 to about 30 % by weight of polyvinylpyrrolidone having an average molecular weight of from about 2,000 to about 54,000; and

d) from about 1 to about 10 % by weight of a surfactant to increase the bioavailability of said ibuprofen.

5 9. The softgel fill formulation of claim 8, wherein said polyethylene glycol is has an average molecular weight of about 600.

10. The softgel formulation of claim 8, wherein said surfactant is selected from esters of d-alpha tocopheryl, polyoxyethylene castor oil derivatives, and polyglycolized glycerides.

10 11. A softgel capsule comprised of a sheath enclosing a liquid fill, said fill comprising:

- a) greater than 30 % by weight ibuprofen in free acid form in solution;
- b) from about 30 to about 60% by weight polyethylene glycol; and
- c) from about 10 to about 30 % by weight of polyvinylpyrrolidone.

15 12. The softgel capsule of claim 11, wherein said ibuprofen is present in an amount of from about 35% to about 40% by weight of said fill.

13. The softgel capsule of claim 11, wherein said polyethylene glycol has an average molecular weight of from about 200 to about 1,000.

20 14. The softgel capsule of claim 11, wherein said surfactant is selected from esters of d-alpha tocopheryl, polyoxyethylene castor oil derivatives, and polyglycolized glycerides.

15. The softgel capsule of claim 11, wherein said polyvinylpyrrolidone has an average molecular weight of from about 2,000 to about 11,000.

16. The softgel capsule of claim 11, wherein said sheath is comprised of from about 45 to about 75% by weight gelatin, from about 20 to about 40 % by weight of a plasticizer, and from about 5 to about 15 % by weight water.

17. The softgel capsule of claim 11, further including up to about 6% of a nasal decongestant or up to about 10% of an antihistaminic.

18. A softgel capsule comprised of a sheath enclosing a liquid fill, said fill comprising:

- 10 a) at least 35 % by weight ibuprofen in free acid form in solution;
- b) from about 30 to about 50% by weight polyethylene glycol having an average molecular weight of from about 200 to about 1,000;
- c) from about 15 to about 30 % by weight of polyvinylpyrrolidone having an average molecular weight of from about 2,000 to about 11,000; and
- 15 d) a surfactant to increase the bioavailability of said ibuprofen.

19. The softgel capsule of claim 18, wherein said surfactant is selected from esters of d-alpha tocopheryl, polyoxyethylene castor oil derivatives, and polyglycolized glycerides.

20. The softgel capsule of claim 18, wherein said sheath is comprised of from about 45 to about 75% by weight gelatin, from about 20 to about 40 % by weight plasticizer, and from about 5 to about 15 % by weight water.

21. The capsule of claim 20, wherein said sheath plasticizer is selected from the group consisting of sorbitol, sorbitol special, maltitol, and mixtures thereof.

22. The capsule of claim 18, wherein the total weight of said capsule fill is less than 600 mg., and the weight of said ibuprofen is at least 200 mg.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/23908

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K31/19 A61K9/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 827 852 A (RUSSELL CARMELITA MACKLIN ET AL) 27 October 1998 (1998-10-27)	1-5, 11-13, 15-17
Y	column 5, line 61 - line 63 column 8; example 3 ---	6-10, 14, 18-22
X	WO 95 19759 A (PROCTER & GAMBLE) 27 July 1995 (1995-07-27) page 14 -page 15; examples 7,8 claims 1-7 --- -/--	1-5, 11-13, 17

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

24 January 2001

Date of mailing of the international search report

05/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Muller, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/23908

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 360 615 A (PATEL MAHENDRA ET AL) 1 November 1994 (1994-11-01) column 4, line 26 -column 5, line 48 column 8, line 51 -column 9, line 29 column 10 -column 11; example 1 column 12; example 9 column 13; example 12 claims 1,5,6,8 ----	1-4, 11-13,15
X	US 5 141 961 A (COAPMAN SCOTT D) 25 August 1992 (1992-08-25) column 3, line 12 -column 9, line 36 claims 1,7,9,11,14,21,24 ----	1,4,5, 11,15-17
X	US 5 641 512 A (CIMILUCA PAUL ALFRED) 24 June 1997 (1997-06-24) the whole document ----	1,4,5, 11,15-17
Y	US 5 431 916 A (WHITE RICHARD K) 11 July 1995 (1995-07-11) column 6, line 13 -column 7, line 11 column 10, line 40 -column 11, line 49; examples 2-6 claims 1,3,5-8 ----	6-10,14, 18-22
P,Y	WO 00 30619 A (ACCUCAPS IND LIMITED) 2 June 2000 (2000-06-02) page 3, line 1 -page 7, line 10 page 8; example 6 claims 1,6,8 -----	6-10,14, 18-22

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/23908

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5827852	A	27-10-1998	CA 2161217 A	10-11-1994
			CN 1122103 A	08-05-1996
			EP 0695175 A	07-02-1996
			FI 955144 A	31-10-1995
			JP 8509725 T	15-10-1996
			NO 954146 A	18-10-1995
			WO 9425009 A	10-11-1994
WO 9519759	A	27-07-1995	AU 706890 B	01-07-1999
			AU 1607595 A	08-08-1995
			BR 9506564 A	02-09-1997
			CA 2181241 A	27-07-1995
			CN 1138827 A	25-12-1996
			CZ 9602104 A	11-12-1996
			EP 0741560 A	13-11-1996
			FI 962948 A	23-07-1996
			HU 74909 A	28-03-1997
			JP 9508128 T	19-08-1997
			NO 963052 A	24-09-1996
			NZ 279443 A	27-04-1998
			PL 315635 A	25-11-1996
			SK 96196 A	05-03-1997
US 5360615	A	01-11-1994	US 5071643 A	10-12-1991
			AT 66810 T	15-09-1991
			AU 606367 B	07-02-1991
			AU 8157387 A	06-05-1988
			CA 1316823 A	27-04-1993
			DE 3772760 A	10-10-1991
			DE 3772760 D	10-10-1991
			EP 0293406 A	07-12-1988
			JP 2564476 B	18-12-1996
			JP 8157354 A	18-06-1996
			JP 2564477 B	18-12-1996
			JP 8157355 A	18-06-1996
			JP 7116021 B	13-12-1995
			JP 1502185 T	03-08-1989
			KR 9406270 B	14-07-1994
			KR 9408030 B	01-09-1994
			KR 9408031 B	01-09-1994
			NZ 222586 A	26-09-1990
			WO 8802625 A	21-04-1988
			ZA 8708723 A	28-06-1989
US 5141961	A	25-08-1992	AU 2191192 A	25-01-1993
			CA 2109623 A,C	07-01-1993
			EP 0591346 A	13-04-1994
			MX 9203447 A	01-12-1992
			PT 100634 A	30-09-1993
			WO 9300072 A	07-01-1993
US 5641512	A	24-06-1997	AU 4988596 A	16-10-1996
			BR 9607961 A	14-07-1998
			CA 2214399 A	03-10-1996
			EP 0817618 A	14-01-1998
			JP 11502838 T	09-03-1999
			WO 9629997 A	03-10-1996

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/23908

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5431916	A	11-07-1995	CA 2159580 A	10-11-1994
			JP 8509498 T	08-10-1996
			WO 9425008 A	10-11-1994
WO 0030619	A	02-06-2000	AU 1172000 A	13-06-2000